

RESEARCH

Open Access



# Comparison of cognitive function, socioeconomic level, and the health-related quality of life between epileptic patients with attention deficit hyperactivity disorder and without

Gellan K. Ahmed<sup>1\*</sup> , Alaa M. Darwish<sup>1</sup>, Hossam Khalifa<sup>1</sup> and Mohamed A. Khashbah<sup>2</sup>

## Abstract

**Background:** Epilepsy is one of the most common neurological conditions. Attention deficit hyperactivity disorder (ADHD) in children with epilepsy proves to be very common. Both epilepsy and ADHD impair quality of life. We aimed to evaluate cognitive function, socioeconomic level, and quality of life (QOL) among children with ADHD and epilepsy. A total of 100 children were divided into 5 groups (20 children/group) as (I) epilepsy, (II) ADHD with epilepsy, (III) ADHD with EEG changes, (IV) ADHD without EEG changes, and (V) control. Children aged between 6 and 11 years were recruited for this study. Early Childhood Epilepsy Severity Scale (E-Chess), Conners' Parent Rating Scale (CPRS), Wechsler Intelligence Scale for Children-3rd edition (WISC-III), socioeconomic scale for assessment of social burden and socioeconomic classes, and PedsQL (quality of life measure) assessed.

**Results:** Children with ADHD and epilepsy had the lowest PedsQL total scores and lower scores than other groups especially in performance IQ score. The highest percentage of low socioeconomic class (25%) was observed in the group of ADHD with epilepsy and the group of epilepsy.

**Conclusion:** ADHD with epilepsy is associated with low performance IQ, poor socioeconomic level, and quality of life. Pediatric Quality of Life Inventory scores show significant correlation with total IQ score in the group of ADHD with epilepsy.

**Keywords:** Epilepsy, ADHD, Quality of life, Socioeconomic, Intelligence

## Background

Epilepsy is the most common neurological condition affecting approximately 65 million people worldwide [1], and it also affects nearly 3.2–5.5/1000 children in the developed world [2].

Epilepsy in children is associated with marked impairment of cognitive functioning [3], increasing the risk of attention deficits and impairing executive functioning [4].

Executive functions are various high-level skills, which allow for flexibility in goal-directed behavior [5], such as response inhibition, working memory, and set-shifting [6]. Additionally, AEDs have been affirmed to increase the risk of cognitive impairment and executive dysfunction in young people with epilepsy [7–9].

Children with epilepsy (CWE) suffer from epileptic seizures and from executive, behavioral, and emotional issues [10]. Deficits in executive functioning are associated with poor health-related quality of life (HRQOL) in these children [11, 12].

\* Correspondence: [gillankaram@aun.edu.eg](mailto:gillankaram@aun.edu.eg)

<sup>1</sup>Department of Neurology and Psychiatry, Assiut University, Assiut 71516, Egypt

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Moreover, the lives of CWE are constantly threatened by physical injuries, such as falls and burns, caused by epilepsy. The fear of acquiring these injuries prevents them from engaging in outdoor activities [13]. They may also need to isolate themselves from society because of the stigma associated with epilepsy [14].

Attention-deficit/hyperactivity disorder (ADHD) is one of the most widespread disorders in CWE [15, 16]. ADHD is generally characterized by persistent inattention, hyperactivity, and impulsivity [17]. Furthermore, several studies found that ADHD children may have electroencephalogram (EEG) abnormalities such as interictal epileptiform discharges (IEDs) or localization of IEDs foci [18, 19]. Lee et al. evaluated the effect of epileptiform discharges in ADHD by comparison between group of ADHD patients with EEG changes and group of ADHD patients with normal EEG. It was found that interictal epileptiform discharges were associated with seizures in children with ADHD [20]. Moreover, a study showed that valproate decreased EEG paroxysm abnormality as well as ADHD scores [21]. This represents pathophysiology overlapping between ADHD and epilepsy [20].

Patients with ADHD suffer from many cognitive deficits because they lack attention and strategic flexibility, struggle to control and inhibit behavior, and have poor planning and working memory capabilities [22–26].

Similar to epilepsy, ADHD is associated with impairment in many aspects of the lives of children who have it, including academic performance, social functioning, and subsequently impaired overall quality of life (QOL) [27–29]. Additionally, ADHD may have a significant economic consequence for children who have it, their families, and society [30].

Hence, epilepsy and ADHD affect the QOL [31]. In the medical researches, “quality of life has become an important measure of outcomes in both research and clinical settings” [32]. There is positive association between QOL impairment and severity of disorder [33]. So, many studies are examining the QOL of patients with epilepsy or ADHD worldwide, particularly in developed countries. However, due to the influence of various variables, such as demographic features and socio-economic and clinical presentations, the QOL of these patients differs from country to country [30].

## Method

### Aim of study

We aim to compare the cognitive function, socioeconomic level, and QOL among epileptic children with ADHD and without.

### Study design

This is a case-control study.

### Site of the study and selection of participants

One hundred children were between 6 and 11 years of age in the period between September 2018 and October 2019. The sample size was calculated with statistical power of 95% by using Epi info 7. Participants were recruited from the Child Psychiatry Clinic of Neurology and Psychiatric Hospital, Assiut University. The children were divided into five groups, namely, as group (1) children with epilepsy alone, group (2) children with ADHD with epilepsy, group (3) children with ADHD with EEG changes, group (4) children with ADHD without EEG changes, and group (5) healthy controls, and each group included 20 children.

We consider using EEG as tool for differing ADHD group as there was evidence that ADHD with EEG changes can be transferred to ADHD with epilepsy after few years [20]. Also, categorizing children with epilepsy by time of seizure onset showed that 15.8% of them were ADHD in early onset of seizures compared to 8.1% in those with newly diagnosed seizures. These results suggested that seizures or subclinical epileptiform bursts play an important role in the development of attention disorders [34]. So, there is bidirectional relationship between both diseases. Our classification to ADHD groups by EEG will help in clarifying this relationship and also, clarify if there is difference between ADHD with EEG changes and without regarding cognitive function and quality of life.

The caregivers of the children and adolescents with epilepsy were interviewed for informed consent and medical information concerning epilepsy and ADHD. ADHD was diagnosed according to DSM-5 [35], and the seizure types were classified according to the International League Against Epilepsy (ILAE) criteria.

### Inclusion criteria

- (i) Diagnosis of ADHD and/or epilepsy was verified and confirmed on a clinical basis.
- (ii) Idiopathic type of epilepsy that was confirmed on clinical basis, neurological assessment, and imaging investigations.

### Exclusion criteria

- (i) Children with significant intellectual disability (Wechsler Intelligence Scale IQ < 70);
- (ii) Children with history or current substance use
- (iii) Children with comorbid medical or other neurological conditions.
- (iv) Children with other psychiatric disorders,

Patients received their medication, as ADHD children received non-stimulant medication (atomoxetine), while epileptic children received their antiepileptic drugs.

### Ethical considerations

This study obtained ethical approval from the Institutional Review Board (IRB) of the Faculty of Medicine, Assiut University, with an approval number of 17200196. This study was also registered on a clinical trial with the following registration number NCT03806946. The parents signed informed consent on behalf of their children to take part in the study.

### Tools

The included children were firstly assessed by semi-structured interview by using the Mini International Neuropsychiatric Interview for Children and Adolescents (MINI KID) Arabic version [36] for psychiatric evaluation then all subjects had the following:

#### *Electroencephalography*

Conventional EEGs were obtained with eight channels and scalp electrodes placed with the international 10–20 system with bipolar and referential montages; it is a Nihon Kohen equipment model. Chloral hydrate was used to induce sleep if needed. Data analysis for EEG was done by determined location and side of none provoked paroxysmal activities, and focal activities if present.

#### *Early Childhood Epilepsy Severity Scale (applied only for the epileptic group) [37]*

The Early Childhood Epilepsy Severity Scale (E-Chess) was developed for use in the Tuberous Sclerosis 2000 Cohort Study to quantify the severity of epilepsy. The scale assesses the severity of seizures by scores on five variables. The variables are frequency of seizures, the time period over which seizures occur, the number of seizure types, the number of antiepileptic drugs used, and response to treatment. Each of these variables was given a score, with a higher score indicating greater severity. Thereafter, the scores were added up to give a total severity score that was used to indicate seizure severity in the current study. E-Chess total score  $\leq 9$  is associated with favorable epilepsy outcomes, while E-Chess total score  $> 10$  is associated with drug-resistant epilepsy with poor responses.

#### *Conners' Parent Rating Scale [38]*

The Conners' Parent Rating Scale (CPRS) is a 110-item parent-rated scale that diagnoses ADHD with other behavioral disorders in children aged 6–18 years. These behavioral disorders are oppositional defiant and conduct disorders.

#### *Wechsler Intelligence Scale for Children-3rd edition (Arabic version by [39])*

Each participant was assessed using the Wechsler Intelligence Scale for Children-3rd edition (WISC-III), which is composed of 13 subtests to test the cognitive abilities of children of different dimensions. The subtest of the WISC includes questions about general knowledge,

traditional arithmetic problems, vocabulary, the completion of mazes, and the arrangement of blocks and pictures. WISC-III provides scores for verbal, performance, and full-scale IQs.

#### *Socioeconomic scale [40]*

The socioeconomic scale is a tool used to assess the social burden and socioeconomic classes. It also contains four main variables: the educational level of the father and the mother, the occupation of the father and the mother, the total family income, and the lifestyle of the family.

#### *Pediatric Quality of Life Inventory, parent-proxy report formats [41]*

The Pediatric Quality of Life Inventory (PedsQL) is a questionnaire assessment tool used to evaluate the QOL of children. It is composed of 23 items that assess functions in the following four areas: physical, emotional, social, and school. The psychosocial health summary score is a computed mean of the emotional, social, and school functioning subscales of the PedsQL. The physical functioning scale is the same as the physical summary score. In addition, the computed mean of the emotional, social, school functioning, and physical scales are used to generate a total summary score. The parents of children reported their response using a 5-point Likert scale ranging from 0 = never to 4 = always, and these responses were associated with a higher score indicating a higher QOL.

### Statistical analysis

The analysis was conducted using SPSS version 26. The frequency and proportion analyses were performed for qualitative variables, while quantitative variables were presented as mean  $\pm$  SD. We used three different statistical tests: the chi-square test was used for analysis between categorical groups, the ANOVA test was employed to detect the presence of significant differences in the mean values for more than two groups, and the Student's *t* test was utilized to test the mean values between two independent groups. Furthermore, the Spearman correlation test was applied for testing correlations between quantitative variables. All the tests were two-tailed, and *p* values of  $< 0.05$  were considered statistically significant.

## Results

### Demographic data

No statistical significance was observed regarding the age of studied groups. Males were significantly higher in all groups ( $p = 0.001$ ), while the group of ADHD with epilepsy comprised 45% of males and 55% of females (see Table 1).

**Table 1** Demographic features of studied group

	Epilepsy (N = 20) N (%)	ADHD with EEG changes (N = 20) N (%)	ADHD without EEG changes (N = 20) N (%)	ADHD with epilepsy (N = 20) N (%)	Control (N = 20) N (%)	P value
Age mean ± SD	8.20 ± 2.462	7.25 ± 1.410	7.33 ± 1.958	7.65 ± 1.348	7.80 ± 1.436	0.448
Sex						0.002*
Male	18 (90%)	18 (90%)	14 (70%)	9 (45%)	12 (60%)	
Female	2 (10%)	2 (10%)	6 (30%)	11 (55%)	8 (40%)	

\*P value is significant

**Clinical features among epilepsy groups**

The clinical features among the group of epilepsy and the group of ADHD with epilepsy validate that a significant difference was observed regarding the types of seizure (*p* value = 0.035) as most of the cases were of generalized type (80% and 100%, respectively). Conversely, no significant differences were detected between the group of epilepsy and the group of ADHD with epilepsy as regards age at the onset of seizure, the duration of epilepsy, or time since the last seizure. However, it was observed that the time since the last seizure between the group of epilepsy and the group of ADHD with epilepsy was less than 1 year (65% and 50%, respectively) (Table 2).

The comparison of the EEG changes among the group of epilepsy, the group of ADHD with epilepsy, and the group of ADHD with EEG changes shows the significant predominance of unprovoked generalized paroxysmal changes in the measured EEG changes (*p* value < 0.000).

Table 3 exhibits that a significant difference exists between the group of epilepsy and the group of ADHD with epilepsy regarding the number of anticonvulsants used and the total E-Chess severity score.

Children who received one or two anticonvulsants showed the highest percentage in the group of epilepsy and the group of ADHD with epilepsy (70%, 80%, respectively).

As regards total E-Chess severity score, children with favorable outcomes (≤ 9) had the highest percentage in the group of epilepsy and the group of ADHD with epilepsy (70% and 100%, respectively).

**Conners 3rd Edition Scale**

All the subscales of the CPRS were significantly different in all groups. The combined type of ADHD represented the highest percentages (70% and 50%, respectively) in the group of ADHD with EEG changes and the group of ADHD with epilepsy. However, the group of ADHD without EEG changes had the highest frequency of hyperactivity type (45%).

Table 4 shows that significant differences among the studied groups in all the CPRS subscales include: inattention, hyperactivity/impulsivity, learning problems, executive functioning, defiance/aggression, and peer/family relation.

The group of ADHD with EEG changes had the worst responses in all the CPRS subscales, where highly elevated clinical ratings represented the highest proportion of all the responses followed by the group of ADHD with epilepsy and ADHD without EEG changes. In contrast, the group of epilepsy had the lowest proportion of clinical ratings in all its subscales, except for the defiance/aggression subscale (55%).

**Wechsler Intelligence Scale for Children-3rd edition**

No significant difference was observed between the studied groups in the IQ classification measured by the WISC-III (*p* value = 0.233).

Moreover, no statistically significant difference emerged among the studied groups in the mean values of verbal and total IQ scores (*p* > 0.05). With regard to performance IQ scores, the group of ADHD with epilepsy had the lowest mean value among all the studied groups (82.28 ± 5.78), while the control group had a significantly high mean value (91.85 ± 6.03) (*p* value = 0.017) (Fig. 1).

The group of ADHD without EEG changes had low performance IQ and total IQ than ADHD with EEG

**Table 2** Clinical features among epilepsy groups

Clinical variables		Epilepsy (N = 20) N (%)	ADHD with epilepsy (N = 20) N (%)	P value
Age of onset of seizure (years)	(means ± SD)	2.35 ± 1.348	2.75 ± 1.209	0.648
Duration of epilepsy (years)	(means ± SD)	5.80 ± 2.142	4.90 ± 1.209	0.084
Time since the last seizure	< 1 year	13 (65%)	10 (50%)	0.107
	1–2 years	7 (35%)	6 (30%)	
	> 2 years	0 (0%)	4 (20%)	
Types of seizure	Partial	4(20%)	0 (0%)	0.035*
	Generalized	16 (80%)	20 (100%)	

\*P value is significant

**Table 3** Early Childhood Epilepsy Severity Scale (E-Chess) among the studied epilepsy groups

Items variables	Epilepsy (N = 20)	ADHD with epilepsy (N = 20)	P value
<b>Frequency of seizures</b>			-
Weekly	20 (100%)	20 (100%)	
Daily	0 (0%)	0 (0%)	
More than daily	0 (0%)	0 (0%)	
<b>Time period over which seizures occurred (months)</b>			-
Less than 1 month	0 (0%)	0 (0%)	
1–6 months	0 (0%)	0 (0%)	
More than 6 months	20 (100%)	20 (100%)	
<b>Number of seizure types</b>			0.244
One seizure type	18 (90%)	20 (100%)	
Two seizure types	2 (10%)	0 (0%)	
Three seizure types	0 (0%)	0 (0%)	
<b>Number of anticonvulsants used</b>			0.006*
None	0 (0%)	4 (20%)	
One or two anticonvulsants	14 (70%)	16 (80%)	
More than two anticonvulsants	6 (30%)	0 (0%)	
<b>Response to treatment</b>			0.052
Complete cessation of seizures	7 (35%)	12 (60%)	
Partial cessation of seizures	13 (65%)	6 (30%)	
No improvement of seizures	0 (0%)	2 (10%)	
<b>Total severity score</b>			
≤ 9	14 (70%)	20 (100%)	0.010*
10–13	6 (30%)	0 (0%)	
13–15	0 (0%)	0 (0%)	
<b>Total severity score (mean ± SD)</b>	9.05 ± 0.759	8.10 ± 0.852	0.001*

\*P value is significant

changes ( $59.01 \pm 16.03$  vs  $61.24 \pm 11.03$  and  $61.64 \pm 16.91$  vs  $63.08 \pm 10.71$ , respectively).

#### Socioeconomic scale

A statistically significant difference emerged between socioeconomic classes among groups. The highest percentage of low socioeconomic class (25%) was observed in the group of ADHD with epilepsy and the group of epilepsy.

Furthermore, a significant difference was observed between the mean values of social class among the studied groups. The control group had a significantly higher mean value of social class scores compared with the studied cases ( $227 \pm 26.350$ ) (Table 5).

The group of ADHD without EEG changes had high socioeconomic scale score relative to the group of ADHD with EEG changes ( $177.72 \pm 64.696$  vs  $148.16 \pm 56.155$ ).

#### Pediatric Quality of Life Inventory

No significant difference was detected between the mean values of the physical summary scores among the studied groups. The group of ADHD with epilepsy had

significantly the worst score of the psychological summary score ( $56.91 \pm 12.98$ ) and the total QOL ( $61.13 \pm 15.96$ ) among all the studied groups (Fig. 2).

The group of ADHD without EEG changes had low psychological summary score and total quality of life score than the group of ADHD with EEG changes ( $59.01 \pm 16.03$  vs  $61.24 \pm 11.03$  and  $61.64 \pm 16.91$  vs  $63.08 \pm 10.71$ , respectively). Meanwhile, the group of ADHD with EEG changes had lower physical summary score than ADHD without EEG changes ( $66.73 \pm 19.05$  vs  $71.08 \pm 21.27$ ).

#### Correlation results


##### Correlation between the total IQ score with socioeconomic scale score and the subscales of PedsQL scores among the studied groups

The socioeconomic score had no significant correlation with the total IQ score in all the studied groups. The physical summary score of PedsQL shows a significant moderate positive correlation with a total IQ score in the group of ADHD with epilepsy. The psychological summary and total QOL scores had a significantly

**Table 4** The Conners' Parent Rating Scale (CPRS) score subscales among the studied groups

Conners' Parent Rating Scale subscales		Epilepsy (N = 20) N (%)	ADHD with EEG changes (N = 20) N (%)	ADHD without EEG changes (N = 20) N (%)	ADHD with epilepsy (N = 20) N (%)	Control (N = 20) N (%)	P value
Inattention	Clinical rating	Very elevated	15 (75%)	8 (40%)	9 (45%)	0 (0%)	< 0.000*
	Borderline	Elevated	0 (0%)	0 (0%)	2 (10%)	0 (0%)	
	Average	High average	4 (20%)	5 (25%)	9 (45%)	0 (0%)	
Hyperactivity/impulsivity	Clinical rating	Very elevated	14 (70%)	7 (35%)	0 (0%)	20 (100%)	< 0.000*
	Borderline	Elevated	14 (70%)	12 (60%)	18 (90%)	0 (0%)	
	Average	High average	3 (15%)	0 (0%)	0 (0%)	0 (0%)	
Learning problems	Clinical rating	Very elevated	17 (85%)	13 (65%)	17 (85%)	1 (5%)	
	Borderline	Elevated	2 (10%)	5 (25%)	2 (10%)	8 (40%)	
	Average	High average	2 (10%)	3 (15%)	0 (0%)	12 (60%)	
Executive functioning	Clinical rating	Very elevated	15 (75%)	9 (45%)	12 (60%)	2 (10%)	< 0.000*
	Borderline	Elevated	0 (0%)	3 (15%)	1 (5%)	0 (0%)	
	Average	High average	1 (5%)	1 (5%)	2 (10%)	3 (15%)	
Defiance/aggression	Clinical rating	Very elevated	19 (95%)	16 (80%)	18 (90%)	1 (5%)	0.002*
	Borderline	Elevated	3 (15%)	1 (5%)	4 (20%)	0 (0%)	
	Average	High average	0 (0%)	5 (25%)	0 (0%)	0 (0%)	
Peer/Family relation	Clinical rating	Very elevated	18 (90%)	16 (80%)	18 (90%)	2 (10%)	< 0.000*
	Borderline	Elevated	1 (5%)	2 (10%)	0 (0%)	2 (10%)	
	Average	High average	0 (0%)	1 (5%)	0 (0%)	5 (25%)	
	Clinical rating	Very elevated	8 (40%)	16 (80%)	18 (90%)	2 (10%)	< 0.000*
	Borderline	Elevated	0 (0%)	3 (15%)	2 (10%)	2 (10%)	
	Average	High average	6 (30)	1 (5%)	0 (0%)	1 (5%)	
	Clinical rating	Very elevated	6 (30)	0 (0%)	0 (0%)	15 (75%)	
	Borderline	Elevated	6 (30)	0 (0%)	0 (0%)	15 (75%)	
	Average	High average	0 (0%)	0 (0%)	0 (0%)	15 (75%)	

\*P value is significant




**Fig. 1** Wechsler Intelligence Scale for Children-3 (WISC-III) score among the studied groups

strong positive correlation with the total IQ score in the group of ADHD with epilepsy, whereas no significant correlations were detected in other groups (see Table 6a and table 6b in supplement data).

**Correlation among socioeconomic score, total IQ score, subscales of PedsQL scores with the clinical variables of epilepsy, and E-Chess total severity score in the group of epilepsy and the group of ADHD with epilepsy**

There was a significantly moderate correlation existed between the age at the onset of seizure and the total IQ score in the group of epilepsy. Meanwhile, other scores did not show a significant correlation (see Table 7 in supplement data).



**Fig. 2** The Pediatric Quality of Life Inventory (PedsQL) scores among the studied groups

**Discussion**

Epilepsy, as a chronic disorder, involves patients in different ways, i.e., physically and socially [42]. Psychiatric-behavioral problems are common in CWE and can have a major influence on their QOL and families. In many cases, these problems are more challenging than epilepsy itself [43]. This study examined cognitive functions, socio-economic levels, and the QOL of ADHD and epilepsy patients.

Although the proportion of males was higher in all groups, except for the group of ADHD with epilepsy, they were almost equal when compared with the controls. Similarly, other studies have reported a prevalence of males with ADHD up to 3–7 times greater than females. Meanwhile, ADHD and epilepsy children often have the same gender ratios [44–46].

In the present study, the relation among epilepsy-related variables showed no significant difference regarding age at the onset of seizure, the duration of epilepsy, and time since the last seizure in CWE-ADHD and

**Table 5** Socioeconomic class scores among the studied groups

Socioeconomic classes	Epilepsy (N = 20) N (%)	ADHD with EEG changes (N = 20) N (%)	ADHD without EEG changes (N = 20) N (%)	ADHD with Epilepsy (N = 20) N (%)	Control (N = 20) N (%)	P value
High classes	0 (0%)	4 (20%)	5 (25%)	0 (0%)	7 (35%)	< 0.000*
Middle classes	15 (75%)	14 (70%)	13 (65%)	15 (75%)	10 (50%)	
Low classes	5 (25%)	2 (10%)	2 (10%)	5 (25%)	3 (15%)	
Total socioeconomic scale score	106.24 ± 30.836	148.16 ± 56.155	177.72 ± 64.696	112.50 ± 35.464	227 ± 26.350	< 0.000*

\*P value is significant

those with epilepsy, while the type of seizure was significantly different between the two groups.

Hermann et al. observed no correlation between younger age at the onset of seizure and the presence of ADHD in the cohort study with a newly diagnosed epilepsy [44], and the same result was found by [47] in a retrospective cohort study on CWE. Contrarily, some studies have demonstrated a significantly high prevalence of ADHD in patients with an earlier epilepsy onset as confirmed by [48]. Moreover, [49] observed that earlier epilepsy onset in younger children was associated with more attention deficit, especially in childhood epilepsy with centrottemporal spikes. ILAE has concluded that the relationship between early seizure onset and ADHD development is still uncertain [50].

Regarding the type of seizures, the higher proportion had a generalized type of epilepsy in both groups. Considerable research has examined whether distinct seizure types were associated with a higher risk of ADHD and epilepsy and has found no association [31, 51, 52]. ILAE has also concluded that specific seizure types are not anticipated higher risk of ADHD in epilepsy [44, 45, 50, 53].

In the current study, the comparison of EEG changes among the three groups, namely, the group of epilepsy, the group of ADHD with epilepsy, and the group of ADHD with EEG changes, showed significant differences for unprovoked generalized paroxysmal changes, which could be attributed to the generalized type of seizures observed in most of the cases.

Additionally, the E-Chess scores in the group of ADHD with epilepsy and the group of epilepsy were significantly different in the number of the anticonvulsants used and the total E-Chess severity score. Regarding the number of the anticonvulsant drugs used, multiple studies have been conducted to clarify the effect of antiepileptic drugs (AEDs) on ADHD comorbidity, but the variety of the anticonvulsant drugs involved restricted the comparison among studies due to their poor consistency. Polytherapy was typically associated with a higher level of behavioral issues according to ILAE [50, 54].

In contrast, several studies from divergent settings have found no correlation between the numbers of anticonvulsant drugs and psychiatric problems [50, 55]. Furthermore, Hermann et al. found no differences between the group of epilepsy and the group of ADHD with epilepsy in terms of the number of anticonvulsant drugs [44]. Regarding seizure frequency, many previous studies have found no correlation of seizure frequency in the studied groups [31, 51, 52].

Regarding the CPRS, a combined subtype of ADHD predominated in the group of ADHD with EEG changes and the group of ADHD with epilepsy, while the hyperactivity subtype predominated in the group of ADHD without EEG changes. Other studies have presented similar results, in which the combined subtype of ADHD

was more prevalent than the inattentiveness subtype in ADHD with epilepsy patients [31, 56–58].

Some studies have corroborated that the combined subtype was more common in patients with ADHD [31, 59]. Meanwhile, other studies have asserted that hyperactivity and impulsivity subtypes were more common [60, 61].

According to EEG, our study categorized ADHD into two subgroups. The group of ADHD with EEG changes showed a predominance of the combined subtype of ADHD similar to ADHD with epilepsy, while the group of ADHD without EEG changes was different from other groups. That was further confirmed by the CPRS subscales, as the group of ADHD with EEG changes similar to the group of ADHD with epilepsy had a highly elevated clinical rating response in all subscales relative to the group of ADHD without EEG changes. These variations in ADHD subtypes could be attributed to the characteristics of the methodology of different studies [51].

In our study, the assessment of cognitive functions by WISC-III questionnaires showed no significant difference in IQ classification among all groups. This is similar to studies that have validated that ADHD patients with and without epilepsy did not vary with respect to IQ ( $p = 0.48$ ), thereby promoting a correlation between epilepsy and ADHD, irrespective of lower IQ scores [57]. Nevertheless, the ADHD and epilepsy groups had lower scores on the measurement of the IQ subscales (verbal, output, and overall IQs) than other groups including the controls with a significant difference in performance IQ.

This was explained by Hermann et al., who argued that motor/psychomotor speed and executive functioning tended to severely affect epilepsy-ADHD children with neuropsychological problems. This was marked by response inhibition impairment, concept formation, mental flexibility/working memory, and passive inattention [44, 56, 62, 63].

According to the socioeconomic scale score, most of the children in the group of ADHD with epilepsy and the group of epilepsy were in low socioeconomic classes with low mean values of social scores in comparison with the other studied groups. This could be explained as epilepsy might be responsible for the decrease of socioeconomic levels.

In a systematic review, there has been growing evidence of the correlation between socio-economic disadvantages and ADHD, indicating that socio-economic disadvantages may lie on the causal pathway between, or may be triggered by, ADHD genotype and phenotype [64].

Several studies reported that low socioeconomic status is a risk factor for the development of epilepsy [65, 66], as low socioeconomic status is associated with social and economic deprivation, unemployment, and low income, which in turn are associated with risk factors as birth defects, trauma, infection, and poor nutrition that may cause epilepsy [67–69].



Another study has revealed that epilepsy prevalence is closely linked to specific socio-economic deprivation measures. When epilepsy is found in areas of greater socioeconomic deprivation, the existing major consequence of epilepsy on employment and higher education may be substantial [70].

Regarding PedsQL, significant differences were detected between groups, while the group of ADHD with epilepsy had the poorest PedsQL scores. The total score and certain subscores of PedsQL (psychological summary score) were the lowest in the group of ADHD with epilepsy. Our findings indicate that the relationship between ADHD and epilepsy co-occurrence with poor PedsQL is greater than that between isolated ADHD or epilepsy and poor PedsQL.

A poor PedsQL may be related to epilepsy, chronic course, social stigma, learning challenges, and seizure fear [71]. Additionally, persistent inattentiveness, hyperactivity, and impulsive ADHD symptoms usually have severe adverse effects on the academic, family, and social life of the children affected [72]. The co-existence of these two conditions may increase the risk of psychosocial and cognitive difficulties and impairment in the QOL.

In the present study, the correlation between the total IQ score with the total socioeconomic scale score and the subscales of PedsQL scores among the studied groups showed that physical summary, psychological summary, and total QOL scores had a significant positive correlation with the total IQ score in the group of ADHD with epilepsy, whereas no significant correlations were detected in other groups.

Similarly, Abd El Naby et al. verified that there was a strong significant correlation between PedsQL scores with IQs ( $p$  value = 0.001) [17]. Other studies have confirmed that ADHD with epilepsy is inversely correlated with the QOL and that treating ADHD significantly improves the QOL of CWE [73, 74].

In our study, the correlation between the total socioeconomic scale, total IQ, subscales of PedsQL scores with the clinical variables of epilepsy and the E-Chess total severity score in the group of epilepsy and the group of ADHD with epilepsy showed a significant correlation between the age at the onset of seizure and the total IQ score in the group of epilepsy. Meanwhile, other scores have not affirmed a significant correlation. In a similar study, young age at the onset of seizure was strongly associated with cognitive impairment in most childhood epilepsies [75].

In contrast, some research has suggested that cognitive dysfunction at the onset of epilepsy does not lead to the increase of intensity over time but keeps on a trajectory [76, 77].

Several factors may have a severe effect on cognitive function in epilepsy, such as underlying structural lesions and epilepsy-related disorders, alternation in neurogenesis and synaptogenesis and change of

excitatory/inhibitory balance, and network connectivity. Other factors such as epileptic activity severity, psychosocial aspects, and surgical or pharmacological treatment of seizures may contribute to increased cognitive impairment in epilepsy [78, 79].

Moreover, no significant correlation emerged between the total E-Chess severity scores with PedsQL, which can be explained as the majority of patients in epilepsy groups had low severity of epilepsy.

Some studies have revealed a positive correlation between higher severity of epilepsy and lower QOL. This could be attributed to the stigma of epilepsy, worries about having seizures in school, cognitive impairment, impaired memory, and inability to pay attention which may be associated with some epilepsy syndromes [80, 81].

Additionally, we found that no significant correlation exists between PedsQL and socioeconomic scale scores with the age at the onset of seizure and the duration of epilepsy. This may be suggested by the prolonged duration of illness that is associated with the adaptation of the child and parents to illness, the acceptance of its stigma with decreased anxiety and depression, and the risk of psychiatric comorbidities that cause the impairment of the QOL.

This study has some limitations. First, our sample size was sufficient for the present study but not large, potentially raising the risk of sample-specific findings and potentially restricting their generalizability to more diverse populations. Second, most of the participants were on medications that had several side effects that should be investigated.

We recommend that all CWE be evaluated for ADHD and given appropriate treatment for them to improve their QOL and protect them against cognitive impairments. Also, further research on a larger group of patients will help support our results.

## Conclusion

Our study demonstrated that ADHD with epilepsy is associated with low performance IQ, poor socioeconomic level, and the QOL. PedsQL shows a significant correlation with the total IQ score.

## Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s43045-020-00054-9>.

**Additional file 1:** Supplemental Data.

## Abbreviations

PedsQL: The Pediatric Quality of Life Inventory; CPRS: Conners' Parent Rating Scale; WISC-III: Wechsler Intelligence Scale for Children-3rd edition; E-Chess: Early Childhood Epilepsy Severity Scale; EEG: Electroencephalography; CWE: Children with epilepsy; ILAE: The International League Against Epilepsy

## Acknowledgements

We would like to thank the Newton-Mosharafa Fund (a collaboration between the Ministry of Higher Education in Egypt and the British Council)

that awarded a scholarship in King's College London with Prof. Patric Bolton. Our deepest thanks for his help and support, step-by-step advisement, and guidance all over this research. The authors would like to thank King's College London for its collaboration.

#### Authors' contributions

GA recruited patients, interpreted the patients' data, and was the contributor in writing the manuscript. AD and HK revised data interpretation, read, and approved the final manuscript. MK helped in data entry, analyzed, and generated result sheets. The author(s) read and approved the final manuscript.

#### Funding

Only Gellan Karamallah was awarded by the Newton–Mosharafa Fund.

#### Availability of data and materials

All data generated or analyzed during this study are available from corresponding author on request.

#### Ethics approval and consent to participate

This study had ethical approval from Institutional Review Board (IRB) of Faculty of Medicine, Assiut University, with approval number 17200196. This study was registered on clinical trial with registration number NCT03806946. The parents signed an informed consent on behalf of their children to take part in the study.

#### Consent for publication

Not applicable

#### Competing interests

The authors declare no conflicts of interests.

#### Author details

<sup>1</sup>Department of Neurology and Psychiatry, Assiut University, Assiut 71516, Egypt. <sup>2</sup>Department of Economics, Neuroeconomics and finance, Claremont Graduate University, Claremont, CA 91711, USA.

Received: 1 July 2020 Accepted: 31 August 2020

Published online: 02 November 2020

#### References

- Moshé SL, Perucca E, Rylvin P, Tomson T (2015) Epilepsy: new advances. *Lancet* (London, England) 385(9971):884–898. [https://doi.org/10.1016/S0140-6736\(14\)60456-6](https://doi.org/10.1016/S0140-6736(14)60456-6)
- Camfield P, Camfield C (2015) Incidence, prevalence and aetiology of seizures and epilepsy in children. *Epileptic Disord* 17(2):117–123. <https://doi.org/10.1684/epd.2015.0736>
- Witt J-A, Helmstaedter C (2017) Cognition in epilepsy: current clinical issues of interest. *Curr Opin Neurol* 30(2):174–179. <https://doi.org/10.1097/WCO.0000000000000430>
- Reuner G, Kadish NE, Doering JH, Balke D, Schubert-Bast S (2016) Attention and executive functions in the early course of pediatric epilepsy. *Epilepsy Behav* 60:42–49. <https://doi.org/10.1016/j.yebeh.2016.04.011>
- Welsh MC, Pennington BF (1988) Assessing frontal lobe functioning in children: views from developmental psychology. *Dev Neuropsychol* 4(3):199–230. <https://doi.org/10.1080/87565648809540405>
- Miyake A, Friedman NP, Emerson MJ, Witzki AH, Howerter A, Wager TD (2000) The unity and diversity of executive functions and their contributions to complex "frontal lobe" tasks: a latent variable analysis. *Cogn Psychol* 41(1):49–100. <https://doi.org/10.1006/cogp.1999.0734>
- Moavero R, Santarone ME, Galasso C, Curatolo P (2017) Cognitive and behavioral effects of new antiepileptic drugs in pediatric epilepsy. *Brain Dev* 39(6):464–469. <https://doi.org/10.1016/j.braindev.2017.01.006>
- Ulate-Campos A, Fernández IS (2017) Cognitive and behavioral comorbidities: an unwanted effect of antiepileptic drugs in children. *Semin Pediatr Neurol* 24(4):320–330. <https://doi.org/10.1016/j.spen.2017.10.011>
- Helmstaedter C, Witt JA, Hoppe C (2019) Evaluating the mediating role of executive functions for antiepileptic drugs' effects on IQ in children and adolescents with epilepsy. *Epilepsy & behavior : E&B* 96:98–103. <https://doi.org/10.1016/j.yebeh.2019.04.003>
- Williams AE, Giust JM, Kronenberger WG, Dunn DW (2016) Epilepsy and attention-deficit hyperactivity disorder: links, risks, and challenges. *Neuropsychiatr Dis Treat* 12:287–296. <https://doi.org/10.2147/NDT.S81549>
- Sherman EM, Slick DJ, Eylr KL (2006) Executive dysfunction is a significant predictor of poor quality of life in children with epilepsy. *Epilepsia*. 47(11):1936–1942. <https://doi.org/10.1111/j.1528-1167.2006.00816.x>
- Love CE, Webbe F, Kim G, Lee KH, Westerveld M, Salinas CM (2016) The role of executive functioning in quality of life in pediatric intractable epilepsy. *Epilepsy Behav* 64(Pt A):37–43. <https://doi.org/10.1016/j.yebeh.2016.08.018>
- Raty LK, Wilde Larsson BM (2007) Quality of life in young adults with uncomplicated epilepsy. *Epilepsy & behavior : E&B* 10(1):142–147. <https://doi.org/10.1016/j.yebeh.2006.09.010>
- Smeets VM, van Lierop BA, Vanhoutvin JP, Aldenkamp AP, Nijhuis FJ (2007) Epilepsy and employment: literature review. *Epilepsy & behavior : E&B* 10(3):354–362. <https://doi.org/10.1016/j.yebeh.2007.02.006>
- Cohen R, Senecky Y, Shuper A, Inbar D, Chodick G, Shalev V et al (2013) Prevalence of epilepsy and attention-deficit hyperactivity (ADHD) disorder: a population-based study. *J Child Neurol* 28(1):120–123. <https://doi.org/10.1177/0883073812440327>
- Besag F, Gobbi G, Caplan R, Sillanpää M, Aldenkamp A, Dunn DW (2016) Psychiatric and behavioural disorders in children with epilepsy (ILAE Task Force Report): epilepsy and ADHD. *Epileptic Disord*. <https://doi.org/10.1684/epd.2016.0811>
- Abd El Naby SA, Naguib YM (2018) Sociodemographic, electrophysiological, and biochemical profiles in children with attention deficit hyperactivity disorder and/or epilepsy. *Behav Neurol* 2018. <https://doi.org/10.1155/2018/8932817>
- Silvestri R, Gagliano A, Calarese T, Arico I, Cedro C, Condruso R et al (2007) Ictal and interictal EEG abnormalities in ADHD children recorded over night by video-polysomnography. *Epilepsy Res* 75(2-3):130–137. <https://doi.org/10.1016/j.eplepsyres.2007.05.007>
- Fonseca LC, Tedrus GMA, Cd M, Machado AV, Almeida MPD, Oliveira DOFd (2008) Epileptiform abnormalities and quantitative EEG in children with attention-deficit/hyperactivity disorder. *Arq Neuropsiquiatr* 66(3A):462–467. <https://doi.org/10.1590/s0004-282x2008000400004>
- Lee EH, Choi YS, Yoon HS, Bahn GH (2015) Clinical impact of epileptiform discharge in children with attention-deficit/hyperactivity disorder (ADHD). *J Child Neurol* 31(5):584–588. <https://doi.org/10.1177/0883073815604223>
- Kanemura H, Sano F, Tando T, Hosaka H, Sugita K, Aihara M (2013) EEG improvements with antiepileptic drug treatment can show a high correlation with behavioral recovery in children with ADHD. *Epilepsy Behav* 27(3):443–448. <https://doi.org/10.1016/j.yebeh.2013.03.014>
- Alloway TP, Passolunghi MC (2011) The relationship between working memory, IQ, and mathematical skills in children. *Learn Individ Differ* 21(1):133–137
- Biederman J, Petty CR, Doyle AE, Spencer T, Henderson CS, Marion B et al (2007) Stability of executive function deficits in girls with ADHD: a prospective longitudinal followup study into adolescence. *Dev Neuropsychol* 33(1):44–61. <https://doi.org/10.1080/87565640701729755>
- Brocki KC, Randall KD, Bohlin G, Kerns KA (2008) Working memory in school-aged children with attention-deficit/hyperactivity disorder combined type: are deficits modality specific and are they independent of impaired inhibitory control? *J Clin Exp Neuropsychol* 30(7):749–759. <https://doi.org/10.1080/13803390701754720>
- Castellanos FX, Tannock R (2002) Neuroscience of attention-deficit/hyperactivity disorder: the search for endophenotypes. *Nat Rev Neurosci* 3(8):617–628. <https://doi.org/10.1038/nrn896>
- Pfiffner LJ, Rooney ME, Jiang Y, Haack LM, Beaulieu A, McBurnett K (2018) Sustained effects of collaborative school-home intervention for attention-deficit/hyperactivity disorder symptoms and impairment. *J Am Acad Child Adolesc Psychiatry* 57(4):245–251. <https://doi.org/10.1016/j.jaac.2018.01.016>
- Bagwell CL, Molina BS, Pelham WE Jr, Hoza B (2001) Attention-deficit hyperactivity disorder and problems in peer relations: predictions from childhood to adolescence. *J Am Acad Child Adolesc Psychiatry* 40(11):1285–1292. <https://doi.org/10.1097/00004583-200111000-00008>
- Sawyer MG, Whaites L, Rey JM, Hazell PL, Graetz BW, Baghurst P (2002) Health-related quality of life of children and adolescents with mental disorders. *J Am Acad Child Adolesc Psychiatry* 41(5):530–537. <https://doi.org/10.1097/00004583-200205000-00010>
- Matza LS, Swensen AR, Flood EM, Secnik K, Leidy NK (2004) Assessment of health-related quality of life in children: a review of conceptual, methodological, and regulatory issues. *Value Health* 7(1):79–92. <https://doi.org/10.1111/j.1524-4733.2004.71273.x>

30. Ashjazadeh N, Yadollahikhales G, Ayoobzadehshirazi A, Sadraei N, Hadi N (2014) Comparison of the health-related quality of life between epileptic patients with partial and generalized seizure. *Iran J Neurol* 13(2):94–100. <https://doi.org/10.1186/1478-7547-3-5>
31. Ekinçi O, Okuyaz Ç, Erdoğan S, Gunes S, Ekinçi N, Kalinlı M et al (2017) Attention-deficit hyperactivity disorder (ADHD) in epilepsy and primary ADHD: differences in symptom dimensions and quality of life. *J Child Neurol* 32(14):1083–1091. <https://doi.org/10.1177/0883073817737445>
32. Agarwal R, Goldenberg M, Perry R, IsHak VVW (2012) The quality of life of adults with attention deficit hyperactivity disorder: a systematic review. *Innov Clin Neurosci* 9(5-6):10–21
33. Danckaerts M, Sonuga-Barke EJS, Banaschewski T, Buitelaar J, Döpfner M, Hollis C et al (2010) The quality of life of children with attention deficit/hyperactivity disorder: a systematic review. *Eur Child Adolesc Psychiatry* 19(2):83–105. <https://doi.org/10.1007/s00787-009-0046-3>
34. Dunn DW, Austin JK, Harezlak J, Ambrosius WT (2003) ADHD and epilepsy in childhood. *Dev Med Child Neurol* 45(1):50–54
35. American Psychiatric Association (2013) Diagnostic and statistical manual of mental disorders teAPA, Washington, DC
36. Ibrahim M, Bishry Z, Hamed A (2002) Comparison of Mini International Neuropsychiatric Interview for children (MINI-KID) with the schedules for affective disorders and schizophrenia for school aged children, present and lifetime version (KSADS-PL). MD thesis. Ain Shams University
37. Humphrey A, Ploubidis GB, Yates JR, Steinberg T, Bolton PF (2008) The early childhood epilepsy severity scale (E-chess). *Epilepsy Res* 79(2-3):139–145. <https://doi.org/10.1016/j.eplepsyres.2008.01.007>
38. Conners KC (2008) Conners 3rd edition. Multi-Health Systems, Toronto
39. Meleka L, Ismail M (1999) Arabic translation of Wechsler Intelligence Scale for children. Egyptian Anglo library, Cairo
40. Abdel-Tawab MA (2010) Socioeconomic scale for family, revised edition. MD thesis in educational basics, Faculty of Education. Assiut University 3:32–55
41. Sabbah I, Sabbah H, Sabbah S, Akoum H, Droubi N, Mercier M (2012) Measurement properties of the Arabic Lebanon version of the Pediatric Quality of Life Inventory 4.0 generic core scales for young child (5-7 years), and child aged 8-12 years: Quality of Life of in Urban and Rural Children in Lebanon. *Creative Educ* 3(06):959
42. Shetty PH, Naik RK, Saroja A, Punith K (2011) Quality of life in patients with epilepsy in India. *J Neurosci Rural Pract* 2(1):33–38. <https://doi.org/10.4103/0976-3147.80092>
43. Besag F, Gobbi G, Aldenkamp A, Caplan R, Dunn DW, Sillanpää M (2016) Psychiatric and behavioural disorders in children with epilepsy (ILAE Task Force Report): behavioural and psychiatric disorders associated with childhood epilepsy syndromes. *Epileptic Disord*. <https://doi.org/10.1684/epd.2016.0815>
44. Hermann B, Jones J, Dabbs K, Allen CA, Sheth R, Fine J et al (2007) The frequency, complications and aetiology of ADHD in new onset paediatric epilepsy. *Brain*. 130(12):3135–3148. <https://doi.org/10.1093/brain/awm227>
45. Hesdorffer DC, Ludvigsson P, Olafsson E, Gudmundsson G, Kjartansson O, Hauser WA (2004) ADHD as a risk factor for incident unprovoked seizures and epilepsy in children. *Arch Gen Psychiatry* 61(7):731–736. <https://doi.org/10.1001/archpsyc.61.7.731>
46. Sherman EM, Slick DJ, Connolly MB, Eylr KL (2007) ADHD, neurological correlates and health-related quality of life in severe pediatric epilepsy. *Epilepsia*. 48(6):1083–1091. <https://doi.org/10.1111/j.1528-1167.2007.01028.x>
47. Kral MC, Lally MD, Boan AD (2016) Identification of ADHD in youth with epilepsy. *J Pediatr Rehabil Med* 9(3):223–229. <https://doi.org/10.3233/PRM-160383>
48. Kwong KL, Lam D, Tsui S, Ngan M, Tsang B, Lam SM (2016) Attention deficit hyperactivity disorder in adolescents with epilepsy. *Pediatr Neurol* 57:56–63. <https://doi.org/10.1016/j.pediatrneurol.2015.12.022>
49. Danhofer P, Pejšochová J, Dušek L, Rektor I, Ošlejšková H (2018) The influence of EEG-detected nocturnal centrotemporal discharges on the expression of core symptoms of ADHD in children with benign childhood epilepsy with centrotemporal spikes (BCECTS): a prospective study in a tertiary referral center. *Epilepsy Behav* 79:75–81. <https://doi.org/10.1016/j.yebeh.2017.11.007>
50. Auvin S, Wirrell E, Donald KA, Berl M, Hartmann H, Valente KD et al (2018) Systematic review of the screening, diagnosis, and management of ADHD in children with epilepsy. Consensus paper of the Task Force on Comorbidities of the ILAE Pediatric Commission. *Epilepsia*. 59(10):1867–1880. <https://doi.org/10.1111/epi.14549>
51. Costa CRCMd, Oliveira GdM, Gomes MdM, Maia Filho HdS (2015) Clinical and neuropsychological assessment of attention and ADHD comorbidity in a sample of children and adolescents with idiopathic epilepsy. *Arq Neuropsiquiatr* 73(2):96–103. <https://doi.org/10.1590/0004-282X20140219>
52. Rheims S, Herbillon V, Villeneuve N, Auvin S, Napuri S, Cancas C et al (2016) ADHD in childhood epilepsy: clinical determinants of severity and of the response to methylphenidate. *Epilepsia*. 57(7):1069–1077. <https://doi.org/10.1111/epi.13420>
53. Dunn DW, Austin JK, Harezlak J (2003) ADHD and epilepsy in childhood. *Dev Med Child Neurol* 45(1):50–54
54. Samaitienė R, Norkūnienė J, Jurkevičienė G, Grikinienė J (2012) Behavioral problems in children with benign childhood epilepsy with centrotemporal spikes treated and untreated with antiepileptic drugs. *Medicina*. 48(7):50
55. Powell K, Walker RW, Rogathe J, Gray WK, Hunter E, Newton CR et al (2015) Cognition and behavior in a prevalent cohort of children with epilepsy in rural northern Tanzania: a three-year follow-up study. *Epilepsy Behav* 51: 117–123. <https://doi.org/10.1016/j.yebeh.2015.06.034>
56. Barkley RA, Murphy KR (2006) Attention-deficit hyperactivity disorder: a clinical workbook. Guilford Press
57. Socanski D, Aurlin D, Herigstad A, Thomsen PH, Larsen TK (2013) Epilepsy in a large cohort of children diagnosed with attention deficit/hyperactivity disorders (ADHD). *Seizure*. 22(8):651–655. <https://doi.org/10.1016/j.seizure.2013.04.021>
58. Graetz BW, Sawyer MG, Hazell PL, Arney F, Baghurst P (2001) Validity of DSM-IV ADHD subtypes in a nationally representative sample of Australian children and adolescents. *J Am Acad Child Adolesc Psychiatry* 40(12):1410–1417. <https://doi.org/10.1097/00004583-200112000-00011>
59. Farah LG, Fayyad JA, Eapen V, Cassir Y, Salamoun MM, Tabet CC et al (2009) ADHD in the Arab world: a review of epidemiologic studies. *J Atten Disord* 13(3):211–222. <https://doi.org/10.1177/1087054708325976>
60. Loutfi KS, Carvalho AM, Lamounier JA, de Almeida Nascimento J (2011) ADHD and epilepsy: contributions from the use of behavioral rating scales to investigate psychiatric comorbidities. *Epilepsy Behav* 20(3):484–489. <https://doi.org/10.1016/j.yebeh.2010.12.043>
61. Zhao Q, Wang M, Kang H, Zhu S (2018) Behavior problems in children with epilepsy and attention-deficit hyperactivity disorder in Central China. *Epilepsy Behav* 89:79–83. <https://doi.org/10.1016/j.yebeh.2018.10.001>
62. Seidman LJ (2006) Neuropsychological functioning in people with ADHD across the lifespan. *Clin Psychol Rev* 26(4):466–485. <https://doi.org/10.1016/j.cpr.2006.01.004>
63. Doyle AE (2006) Executive functions in attention-deficit/hyperactivity disorder. *J Clin Psychiatry* 67:21–26
64. Russell AE, Ford T, Williams R, Russell G (2016) The association between socioeconomic disadvantage and attention deficit/hyperactivity disorder (ADHD): A Systematic Review. *Child Psychiatry Hum Dev* 47(3):440–458. <https://doi.org/10.1007/s10578-015-0578-3>
65. Hesdorffer DC, Tian H, Anand K, Hauser WA, Ludvigsson P, Olafsson E et al (2005) Socioeconomic status is a risk factor for epilepsy in Icelandic adults but not in children. *Epilepsia*. 46(8):1297–1303. <https://doi.org/10.1111/j.1528-1167.2005.10705.x>
66. Heaney DC, MacDonald BK, Everitt A, Stevenson S, Leonardi GS, Wilkinson P et al (2002) Socioeconomic variation in incidence of epilepsy: prospective community based study in south east England. *BMJ (Clinical research ed)* 325(7371):1013–1016. <https://doi.org/10.1136/bmj.325.7371.1013>
67. Winkley MA, Jatulis DE, Frank E, Fortmann SP (1992) Socioeconomic status and health: how education, income, and occupation contribute to risk factors for cardiovascular disease. *Am J Public Health* 82(6):816–820. <https://doi.org/10.2105/ajph.82.6.816>
68. Karp A, Käreholt I, Qiu C, Bellander T, Winblad B, Fratiglioni L (2004) Relation of education and occupation-based socioeconomic status to incident Alzheimer's disease. *Am J Epidemiol* 159(2):175–183. <https://doi.org/10.1093/aje/kwh018>
69. Cubbin C, LeClere FB, Smith GS (2000) Socioeconomic status and the occurrence of fatal and nonfatal injury in the United States. *Am J Public Health* 90(1):70–77. <https://doi.org/10.2105/ajph.90.1.70>
70. Steer S, Pickrell WO, Kerr MP, Thomas RH (2014) Epilepsy prevalence and socioeconomic deprivation in England. *Epilepsia*. 55(10):1634–1641. <https://doi.org/10.1111/epi.12763>
71. Giuliano L, Cicero CE, Padilla S, Mayaregua DR, Villarreal WMC, Sofia V et al (2019) Knowledge, stigma, and quality of life in epilepsy: results before and after a community-based epilepsy awareness program in rural Bolivia. *Epilepsy Behav* 92:90–97. <https://doi.org/10.1016/j.yebeh.2018.11.036>

72. Haack LM, Villodas M, McBurnett K, Hinshaw S, Pfiffner LJ (2017) Parenting as a mechanism of change in psychosocial treatment for youth with ADHD, predominantly inattentive presentation. *J Abnorm Child Psychol* 45(5):841–855. <https://doi.org/10.1007/s10802-016-0199-8>
73. Yoo HK, Park S, Wang H-R, Lee JS, Kim K, Paik K-W et al (2009) Effect of methylphenidate on the quality of life in children with epilepsy and attention deficit hyperactivity disorder. *Epileptic Disorders* 11(4):301–308. <https://doi.org/10.1684/epd.2009.0278>
74. Radziuk AL, Kieling RR, Santos K, Rotert R, Bastos F, Palmieri AL (2015) Methylphenidate improves the quality of life of children and adolescents with ADHD and difficult-to-treat epilepsies. *Epilepsy Behav* 46:215–220. <https://doi.org/10.1016/j.yebeh.2015.02.019>
75. Hermann B, Jones J, Sheth R, Dow C, Koehn M, Seidenberg M (2006) Children with new-onset epilepsy: neuropsychological status and brain structure. *Brain*. 129(10):2609–2619. <https://doi.org/10.1093/brain/awl196>
76. Baillet LL, Turk WR (2000) The impact of childhood epilepsy on neurocognitive and behavioral performance: a prospective longitudinal study. *Epilepsia*. 41(4): 426–431. <https://doi.org/10.1111/j.1528-1157.2000.tb00184x>
77. Hermann BP, Jones JE, Sheth R, Koehn M, Becker T, Fine J et al (2008) Growing up with epilepsy: a two-year investigation of cognitive development in children with new onset epilepsy. *Epilepsia*. 49(11):1847–1858. <https://doi.org/10.1111/j.1528-1167.2008.01735.x>
78. Berg AT, Caplan R, Hesdorffer DC (2011) Psychiatric and neurodevelopmental disorders in childhood-onset epilepsy. *Epilepsy Behav* 20(3):550–555. <https://doi.org/10.1016/j.yebeh.2010.12.038>
79. Holmes GL (2016) Effect of seizures on the developing brain and cognition. *Seminars in pediatric neurology*. Elsevier, pp 120–126
80. Sabaz M, Cairns DR, Lawson JA, Nheu N, Bleasel AF, Bye AM (2000) Validation of a new quality of life measure for children with epilepsy. *Epilepsia*. 41(6):765–774. <https://doi.org/10.1111/j.1528-1157.2000.tb00240.x>
81. Clary LE, Vander Wal JS, Titus JB (2010) Examining health-related quality of life, adaptive skills, and psychological functioning in children and adolescents with epilepsy presenting for a neuropsychological evaluation. *Epilepsy Behav* 19(3):487–493. <https://doi.org/10.1016/j.yebeh.2010.08.002>

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)

---